



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

MECHANICS.

Conducted by B. F. FINKEL, Kidder, Mo. All contributions to this department should be sent to him.

SOLUTIONS TO PROBLEMS.

9. Proposed by CHARLES E. MYERS, Canton, Ohio.

A ladder inclined at an angle of 60° to the horizon rests with one end on a rough pavement and the other against a smooth vertical wall. If the coefficient of friction between the foot of the ladder and the pavement is $\frac{1}{3}$, to what height can a man ascend before the ladder will begin to slip?

Solution by B. F. FINKEL, Professor of Mathematics, Physics, and Astronomy, Kidder Institute, Kidder, Missouri.

Let AB be the ladder, length $2l$; θ its inclination to the horizon, W its weight and G its center of gravity; $AP=x$, the distance the man ascends and w his weight; I , the center of gravity of the man and ladder; $F=\mu$, the coefficient of friction; and S and R the normal resistance at A and B , respectively.

Then we have $S-F=0 \dots (1)$ and, $W+w-R=0 \dots (2)$. Taking moments about I , we have $W \cdot GI = w \cdot PI$, or $W \cdot GI - w \cdot PI = 0 \dots (3)$. But $GI + PI = x - \frac{1}{2}l$. Hence, from (3), GI

$= \frac{w}{W+w} (x - \frac{1}{2}l)$. Taking moments about A ,

$(W+w) \cdot AI = AB \cdot R(\cos \theta - \mu \sin \theta)$ or $(W+w)$

$\left[\frac{w}{W+w} (x - \frac{1}{2}l) + \frac{1}{2}l \right] \cos \theta = 2lR(\cos \theta - \mu \sin \theta) \dots (4)$. (2) and (4) give

$2l(W+w)(\cos \theta - \mu \sin \theta) = (Wl + wx) \cos \theta$, whence

$x = \frac{2l(W+w)(1 - \mu \tan \theta - W)}{w}$. But $\mu = \frac{1}{3}$ and $\tan \theta = \sqrt{3}$.

$\therefore AP = x = l$. Hence the man can ascend to the middle point of the ladder.

[Note.—This problem and its solution was selected for publication in the Aug. No. But it was lost in the office of the publishers and thus we are unable to credit our contributors for their solutions to it. ED.]

10. Proposed by G. B. M. ZERR, A. M., Principal of High School, Staunton Virginia.

A paraboloid floats in a liquid which fills a fixed paraboloid shell; both the paraboloid and the shell have their axis vertical and their vertices downward; the latus rectum of the paraboloid and shell are equal, and the axis of the shell is m times that of the paraboloid. If the paraboloid be pressed down until its vertex reaches the vertex of the shell, so that some of the liquid overflows, and then released, it is found that the paraboloid rises until it is just wholly out of the liquid, and then begins to fall. Prove that (1) the densities of the paraboloid and liquid are in the ratio.

$2[m^2 + m + 1 = (m + 1)(\sqrt{m^2 - 1})] : 3\sqrt{(m + 1) + (m - 1)}$, the free surface of the liquid being supposed to remain horizontal throughout the motion; and (2)

